FORM B.

Return showing the description and weight of all yarn spun in the Cotton Mills in the Mysore State during the month ending 31st August 1940.

Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count
2s	27,877	16s	44,801	42s	12,204
4s	8,873	16½s	7,321	44s	10,159
6s	6,324	18s	3,952	50s	2,063
6½s	47,138	19s	62,385	60s	34
8s	29,943	20s	332,876		
8½s	110,858	20½s	19,547		
9s	1,414	21s	151,011		
10s	1,034	22s	97,842		
10½s	231,097	24s	115,671		
11s	206,681	26s	33,976		
12s	33,119	27	296		
12½s	83,143	28s	27,380		
13s	1,893	30s	49,468		
13½s	5,716	32s	36,326		
14s	316,803	36s	47,338		
14½s	41,733	38s	2,703		
15s	36,079	40s	62,053	Total ...	2,309,131

BANGALORE,
Dated 28th September 1940.

V. VENKATACHAR,
For *Director*.

MYSORE STATE DEPARTMENT OF PUBLIC HEALTH.

NOTE ON THE STATE OF PUBLIC HEALTH IN MYSORE STATE FOR THE WEEK ENDED THURSDAY, OCTOBER 3, 1940.

A total of 65 attacks with 39 deaths from Plague was returned during the week against 56 attacks with 33 deaths in the previous week. Bangalore City reported nil attack with 1 death while Mysore City and Kolar Gold Field Area continued to remain free from the disease. Comparative figures for the corresponding weeks in 1939 and 1938 were 31 attacks with 19 deaths and 34 attacks with 23 deaths, respectively.

A total of 1 attack with nil death from small-pox was reported from a village of French-Rocks Taluk.

The State remained free from Cholera, Influenza and Relapsing Fever.

The total mortality for Bangalore City amounted to 129 (55 in hospitals), representing a calculated death-rate of 30·1 (17·3 excluding hospital deaths) per mille of population as against 28·0 for the corresponding week of last year. Classified according to causes, the deaths numbered 1 under Plague, 3 under Malaria, 2 under Typhoid, 5 under other Fevers, 5 under Phthisis, 27 under Dysentery and Diarrhœa, and 86 under all other causes. The reported mortality was high due to increase in deaths under Dysentery and Diarrhœa.

During the week, the Civil and Military Station, Bangalore, returned 79 deaths (12 in hospitals), representing a calculated death-rate of 27·7 (23·8 excluding hospital deaths) per mille of population as against 23·7 for the corresponding week of last year. Classified according to causes, the deaths were Broncho-Pneumonia 2, Pneumonia 8, Typhoid 1, Phthisis 3, Dysentery and Diarrhœa 16, and all other causes 49. The reported mortality was high due to increase in deaths under Dysentery and Diarrhœa.

The total mortality for Mysore City was 43 for 6 days (15 in hospitals), representing a calculated death-rate of 20·3 (13·2 excluding hospital deaths) per mille of population as against 27·9 for the corresponding week of last year. Classified according to causes, the deaths were : Other Fevers 3, Phthisis 6, Dysentery and Diarrhœa 6 and all other causes 28. The reported mortality was very low.

There were 54 deaths (7 in hospitals) from all causes in Kolar Gold Field Area equivalent to a calculated death-rate of 34·0 (29·6 excluding hospital deaths) per mille of population as against 29·5 for the corresponding week of last year. Classified according to causes, the deaths numbered—1 under Malaria, 1 under Typhoid, 8 under other Fevers, 2 under Phthisis, 6 under Dysentery and Diarrhœa and 36 under all other causes. The reported mortality was normal.

E. R. SUNDARARAJAN,
For *Director*.